


Technical Information about Rp-2'-AHC-cAMPS-Agarose

Gel for affinity chromatography of cAMP-dependent protein kinases and phosphodiesterases

Update: September 07, 2018 HJ



Abbreviation: Rp-2'-AHC-cAMPS-Agarose

BIOLOG Cat. No.: A 049

Description: In Rp-2'-AHC-cAMPS-Agarose the Rp-isomer of 2'-(6-aminohexylcarbamoyl)adenosine-3', 5'-cyclic monophosphorothioate has been immobilized as an affinity ligand by the hydroxysuccinimide technique. Rp-2'-AHC-cAMPS (Cat. No. A 120) is an analogue of the parent second messenger cyclic AMP in which one of the two exocyclic oxygen atoms in the cyclic phosphate moiety is modified by sulfur. Equatorial thio substitution leads to the R-isomer, while axial modification yields the corresponding S-compound. The suffix "p" indicates that R/S nomenclature refers to phosphorus. The 2'-hydroxy group carries the aminohexylcarbamoyl spacer for attachment.

Properties: Rp-2'-AHC-cAMPS-Agarose is a gel for affinity chromatography of various cAMP-responsive proteins, especially those which tolerate modification of the 2'-ribose hydroxyl group, such as certain phosphodiesterases. Due to its phosphorothioate modification the affinity to most cAMP receptor proteins are weaker, so desorption procedure should be milder. In addition, destruction of the ligand by phosphodiesterases is blocked.

Related Products: This type of gel is also available with the corresponding Sp-isomer (Sp-2'-AHC-cAMPS-Agarose, Cat. No. A 050) and the normal, sulfur-free ligand (2'-AHC-cAMP-Agarose, Cat. No. A 058). For immobilized cGMP or cAMP with other spacer types, attachment positions, different immobilization techniques or on other gels please inquire.

Specification: Ligand density: approximately 6 $\mu\text{mol/ml}$ of settled gel. UV: λ_{max} 258 nm/suspension in glycol

Stability and Storage: Rp-2'-AHC-cAMPS-Agarose has sufficient stability for chromatography at ambient temperature and does not need special care during handling or shipment. Nevertheless, for longer storing periods the gel should be kept in the refrigerator at +4 - +8 °C. **Storage buffer should contain 0.1 % sodium azide for prevention of microbial growth.**

Chromatography: After equilibration with about 10 column volumes of starting buffer the affinity column is loaded with the protein solution, e.g. at 50-200 $\mu\text{l/min}$. In order to elute other nucleotide-dependent proteins unspecifically bound, the column is washed, e.g. with 1 mM 5'-AMP. For elution of E. coli nucleoside diphosphate kinase it is recommended to wash the column with 10 mM ADP/20mM MgCl_2 .

Elution of the target protein is performed by a cyclic nucleotide gradient up to 40 mM. For elution of phosphodiesterases it could be advisable to use the highly hydrolysis-resistant phosphorothioate-modified analogs or PDE inhibitors.

Suitable buffer systems for your special application have to be tested, but phosphate should be not optimal since one essential affinity interaction of cyclic nucleotides towards their target receptors is the cyclic phosphate.

Regeneration can be achieved by a combination of up to 100 mM cAMP and buffer salts or 8 M urea.

Selected References for cAMP Affinity Chromatography:

Van Bemmelen, M.X.P, Ph.D. Thesis, University of Bremen, 1993, Bremen, Germany

Iwitzki, F.; Van Bemmelen, M.X.P.; Genieser, H.-G.; Jastorff, B., Proceedings of the 3rd Swedish-German Workshop on Nucleic Acid Synthesis, Structure and Function, Uppsala, Sweden, 1992: "cAMP Analogues Designed for Affinity Chromatography"

Iwitzki, F.; Van Bemmelen, M.X.P.; Genieser, H.-G.; Jastorff, B., Proc. Internatl. Conf. Second Messengers and Phosphoprot., Glasgow, UK 1992: "New Affinity Materials for Cyclic AMP Binding Proteins"

Dills, W.L.; Beavo, J.A.; Bechtel, P.J.; Myers, K.R.; Sakai, L.J., Krebs, E.G., *Biochemistry* **15**, 3724 - 3770 (1976): "Binding of Adenosine 3',5'-Monophosphate Dependent Protein Kinase Regulatory Subunit to Immobilized Cyclic Nucleotide Derivatives"

Copyright September 18 by BIOLOG Life Science Institute

Dills, W.L.; Beavo, J.A.; Bechtel, P.J.; Krebs, E.G., *Biochem. Biophys. Res. Commun.*, **62**, 70 - 77 (1975): "Purification of Rabbit Skeletal Muscle Protein Kinase Regulatory Subunit Using Cyclic Adenosine- 3':5'- Monophosphate Affinity Chromatography"