


Technical Information about 2'- / 3'- O- (2- Aminoethyl-carbamoyl)- ATP (2'-/3'-AEC-ATP / EDA-ATP)

Update: November 01, 2018 HU



Abbreviation:

2'-/3'-AEC-ATP / EDA-ATP

Formula	CAS No.	Molecular Weight	UV	BIOLOG Cat. No.
C ₁₃ H ₂₂ N ₇ O ₁₄ P ₃ for free acid	[173074-70-9]	593.3 for free acid	λ _{max} 259 nm / ε 15000 / pH 7	A 072

Name: 2'- / 3'- O- (2- Aminoethylcarbamoyl)adenosine- 5'- O- triphosphate

Description: 2'-/3'-AEC-ATP is an analogue of the natural structure adenosine- 5'- O- triphosphate (ATP) in which an ethyl spacer with a terminal amino group has been attached either to the ribose 2'-hydroxy group or to the 3'-hydroxy group by a carbamate bond.

Properties:

- ATP analogue suitable for immobilization as a ligand for affinity chromatography,
- useful for coupling of various labelling structures including fluorophores.

Specification: Sodium salt in aqueous solution (10 mM). The free acid or other salt forms are available upon request. Micro molar quantities are determined by UV at λ_{max}. When opening the tube please make sure that no liquid is lost within the cap. A short spin-down in a bench centrifuge is recommended before use.

Purity: Typical analysis is better than 95% (HPLC / UV / 259 nm) at time of quality control and packing. The product is not sterile and has not been tested for endotoxins.

Stability and Storage: 2'-/3'-AEC-ATP is relatively stable when stored frozen in aqueous solution (-20° celsius necessary, -80° recommended). In order to maintain its original high quality, it is recommended to allow thawing only before using the product. If you will not use up the vial with one application, please aliquot the content of the vial in order to avoid repeated freeze/thaw cycles for the rest. When making such aliquots be sure to operate quickly and to freeze the vial again as soon as possible.

Toxicity and Safety: Since nucleoside triphosphates have multiple tasks in every organism, it is very likely that ATP analogues will interfere with many cell regulation processes *in vivo*. However, due to the rather small quantities to work with, no health hazards have been reported. Nevertheless please keep in mind, that the *in vivo* properties of this compound are not sufficiently characterized up to now. Avoid skin contact or ingestion and allow only trained personnel to handle the product.

Our products are designed, developed and sold for research purposes only! They are intended for *in vitro* and nonhuman *in vivo* laboratory applications. Any other use requires approval of health authorities.

Not for drug, household or related uses!

Selected References for 2'-/3'-AEC-ATP / EDA-ATP:

Webb, M.R.; Corrie, J.E., *Biophys. J.*, **81**, 1562 - 1569 (2001): "Fluorescent Coumarin-labeled Nucleotides to Measure ADP Release from Actomyosin"

Oiwa, K.; Eccleston, J.F.; Anson, M.; Kikumoto, M.; Davis, C.T.; Reid, G.P.; Ferenczi, M.A.; Corrie, J.E.T.; Yamada, A.; Nakayama, H.; Trentham, D.R., *Biophys. J.*, **78**, 3048 - 3071 (2000): "Comparative Single-Molecule and Ensemble Myosin Enzymology: Sulfoindocyanine ATP and ADP Derivatives"

Oiwa, K.; Yamaga, T.; Yamada, A., *J. Biochem.*, **123**, 614 - 618 (1998): "Direct Observation of a Central Bare Zone in a Native Thick Filament Isolated from the Anterior Byssus Retractor Muscle of *Mytilus edulis* Using Fluorescent ATP Analogue"