


Technical Information about c[8-Biotin-11-G(2',5')pA(3',5')p]

Update: August 06, 2019 HU



Abbreviation: c[8-Biotin-11-G(2',5')pA(3',5')p] / c[8-[Biotin]-AET-G(2',5')pA(3',5')p]

Formula	CAS No.	Molecular Weight	UV	BIOLOG Cat. No.
C ₃₈ H ₅₄ N ₁₄ O ₁₆ P ₂ S ₂ (free acid)	[pending]	1089.0 (free acid)	λ _{max} 266 nm / ε 23500 / pH 7	C 176

Name: Cyclic (8- (2- (6- [biotinyl]amino)hexanoyl)- aminoethylthio)guanosine- (2' → 5')- monophosphate- adenosine- (3' → 5')- monophosphate)

Syn.: 8-[Biotin]-AET-cGAMP(2'-5') / 8-[Biotin]-AET-2'3'-cGAMP / 8-[Biotin]-AET-2',5'-3',5'-cGAMP

Description: c[8-Biotin-11-G(2',5')pA(3',5')p] is a biotinylated analogue of the metazoan cyclic dinucleotide second messenger c[G(2',5')pA(3',5')p] (*aka* cGAMP(2'-5') or 2'3'-cGAMP, Cat. No. C 161) in which the biotin label is attached to position 8 of the guanine nucleobase via a 11 atom spacer.

Properties: Analogue of c[G(2',5')pA(3',5')p] with biotin conjugate most probably suitable as tracer in immunoassays. c[G(2',5')pA(3',5')p] was found to be the metazoan second messenger produced by the mammalian innate immune DNA sensor cGAMP synthase (cGAS).

Specification: Crystallized or lyophilized sodium salt. Please keep in mind that equal concentrations of the compound may look different in volume due to sensitivity of the lyophilized form to humidity. The compound can even contract to small volume droplets. Normally the product is located in the conical bottom of the tube. Micromolar quantities are determined by UV at λ_{max}.

Purity: Typical analysis is better than 95% (HPLC / UV / 266 nm). The product is not sterile and has not been tested for endotoxins.

Solubility: c[8-Biotin-11-G(2',5')pA(3',5')p] is soluble in water and aqueous buffers (> 1.5 mM, limits have not been determined). Please rinse tube walls carefully and preferably use ultrasonic or vortex to achieve total and uniform mixing. When opening the tube please make sure that no substance is lost within the cap.

Stability and Storage: c[8-Biotin-11-G(2',5')pA(3',5')p] has sufficient stability at room temperature and does not need special care during handling or shipment. Nevertheless, we recommend that the compound should be stored in the freezer, for longer storage periods preferably in freeze-dried form.

Toxicity and Safety: Please keep in mind, that the *in vivo* properties of this compound are not sufficiently characterized up to now. Avoid skin contact or ingestion and allow only trained personnel to handle the product.

Our products are designed, developed and sold for research purposes only! They are intended for *in vitro* and nonhuman *in vivo* laboratory applications. Any other use requires approval of health authorities.

Not for drug, household or related uses!

Selected References for c[8-Biotin-11-G(2',5')pA(3',5')p]:

Bridgeman, A.; Maelfait, J.; Davenne, T.; Partridge, T.; Peng, Y.; Mayer, A.; Dong, T.; Kaever, V.; Borrow, P.; Rehwinkel, J., *Science*, **349**, 1228 - 1232 (2015): "Viruses Transfer the Antiviral Second Messenger cGAMP Between Cells"

Selected References for the Parent Compound c[G(2',5')pA(3',5')p] (Cat. No. C 161):

Lolicato, M.; Bucchi, A.; Arrigoni, C.; Zucca, S.; Nardini, M.; Schroeder, I.; Simmons, K.; Aquila, M.; DiFrancesco, D.; Bolognesi, M.; Schwede, F.; Kashin, D.; Fishwick, C.W.; Johnson, A.P.; Thiel, G.; Moroni, A., *Nat. Chem. Biol.*, **10**, 457 - 462 (2014): "Cyclic Dinucleotides bind the C-linker of HCN4 to Control Channel cAMP Responsiveness"

Yi, G.; Brendel, V.P.; Shu, C.; Li, P.; Palanathan, S.; Kao, C.C., *PLoS One*, **8**, e77846 (2013): "Single Nucleotide Polymorphisms of Human STING Can Affect Innate Immune Response to Cyclic Dinucleotides"

Konno, H.; Konno, K.; Barber, G.N., *Cell*, **155**, 688 - 698 (2013): "Cyclic Dinucleotides Trigger ULK1 (ATG1) Phosphorylation of STING to Prevent Sustained Innate Immune Signaling"

Ablasser, A.; Schmid-Burgk, J.L.; Hemmerling, I.; Horvath, G.L.; Schmidt, T.; Latz, E.; Hornung, V., *Nature*, **503**, 530 - 534 (2013): "Cell Intrinsic Immunity Spreads to Bystander Cells via the Intercellular Transfer of cGAMP"

Li, X.D.; Wu, J.; Gao, D.; Wang, H.; Sun, L.; Chen, Z.J., *Science*, **341**, 1390 - 1394 (2013): "Pivotal Roles of cGAS-cGAMP Signaling in Antiviral Defense and Immune Adjuvant Effects"

Gao, D.; Wu, J.; Wu, Y.T.; Du, F.; Aroh, C.; Yan, N.; Sun, L.; Chen, Z.J., *Science*, **341**, 903 - 906 (2013): "Cyclic GMP-AMP Synthase is an Innate Immune Sensor of HIV and Other Retroviruses"

Gao, P.; Ascano, M.; Wu, Y.; Barchet, W.; Gaffney, B.L.; Zillinger, T.; Serganov, A.A.; Liu, Y.; Jones, R.A.; Hartmann, G.; Tuschl, T.; Patel, D.J., *Cell*, **153**, 1094 - 1107 (2013): "Cyclic [G(2',5')pA(3',5')p] Is the Metazoan Second Messenger Produced by DNA-Activated Cyclic GMP-AMP Synthase"

Ablasser, A.; Goldeck, M.; Cavlar, T.; Deimling, T.; Witte, G.; Röhl, I.; Hopfner, K.-P.; Ludwig, J.; Hornung, V., *Nature*, **498**, 380 - 384 (2013): "cGAS Produces a 2'-5'-linked Cyclic Dinucleotide Second Messenger that Activates STING"

Zhang, X.; Shi, H.; Wu, J.; Zhang, X.; Sun, L.; Chen, C.; Chen, Z.J., *Mol. Cell*, **51**, 226 - 235 (2013): "Cyclic GMP-AMP Containing Mixed Phosphodiester Linkages Is An Endogenous High-Affinity Ligand for STING"

Diner, E.J.; Burdette, D.L.; Wilson, S.C.; Monroe, K.M.; Kellenberger, C.A.; Hyodo, M.; Hayakawa, Y.; Hammond, M.C.; Vance, R.E., *Cell Rep.*, **3**, 1355 - 1361 (2013): "The Innate Immune DNA Sensor cGAS Produces a Noncanonical Cyclic Dinucleotide that Activates Human STING"

Wu, J.; Sun, L.; Chen, X.; Du, F.; Shi, H.; Chen, C.; Chen, Z.J., *Science*, **339**, 826 - 830 (2013): "Cyclic GMP-AMP is an Endogenous Second Messenger in Innate Immune Signaling by Cytosolic DNA"

Sun, L.; Wu, J.; Du, F.; Chen, X.; Chen, Z.J., *Science*, **339**, 786 - 791 (2013): "Cyclic GMP-AMP Synthase is a Cytosolic DNA Sensor That Activates the Type I Interferon Pathway"