


Technical Information about 8-NO₂-cGMP

Update: May 10, 2022 ss



Abbreviation:

8-NO₂-cGMP

Formula	CAS No.	Molecular Weight	UV	BIOLOG Cat. No.
C ₁₀ H ₁₀ N ₆ O ₉ P · Na	[913645-39-3]	412.2	λ _{max} 396 nm / ε 4100 / pH 7	N 011

Name: 8- Nitroguanosine- 3', 5'- cyclic monophosphate, sodium salt

Description: 8-NO₂-cGMP is an analogue of the parent compound cyclic GMP (cGMP, Biolog Cat. No. G 001) in which the hydrogen in position 8 of the heterocyclic nucleobase is replaced by NO₂.

Properties: The nitrated derivative of cGMP, 8-NO₂-cGMP (8-Nitro-cGMP), is considered to play an important role in NO-mediated signal transduction. Formation of protein Cys-cGMP adducts by 8-NO₂-cGMP was identified as a post-translational modification of proteins, which is called S-guanylation.

Specification: Lyophilized or crystallized solid. Equal concentrations of 8-NO₂-cGMP can appear very different in volume due to sensitivity of the lyophilized form to humidity. The compound can even contract to small volume droplets. Normally the product is located in the conical bottom of the tube. Micromolar quantities are determined by UV at λ_{max}.

Purity: Typical analysis is better than 98 % (HPLC / UV / 396 nm). The product is not sterile and has not been tested for endotoxins.

Solubility: 8-NO₂-cGMP is soluble in water (≥ 39 mM, limits have not been determined). Please rinse tube walls carefully and preferably use ultrasonic or vortex to achieve total and uniform mixing. When opening the tube please make sure that no substance is lost within the cap.

Stability and Storage: 8-NO₂-cGMP has sufficient stability for short-term exposure to room temperature and does not need special care during handling or shipment. Nevertheless, the compound should be protected from light and stored in the freezer, for longer storage periods preferably in freeze-dried form.

Toxicity and Safety: Please keep in mind that the *in vivo* properties of this compound are not sufficiently characterized up to now. Avoid skin contact or ingestion and allow only trained personnel to handle the product.

Our products are designed, developed and sold for research purposes only! They are intended for *in vitro* and nonhuman *in vivo* laboratory applications. Any other use requires approval of health authorities.

Not for drug, household or related uses!

Selected References for 8-NO₂-cGMP:

Petrivalský, M.; Luhová, L.; *Front. Plant Sci.*, **11**:598 (2020): "Nitrated Nucleotides: New Players in Signaling Pathways of Reactive Nitrogen and Oxygen Species in Plants"

Tsutsuki, H.; Kasamatsu, S.; Kunieda, K.; Ida, T.; Sawa, T.; Sasakawa, N.; Akaike, T.; Ihara, H., *Biochem. Biophys. Res. Commun.*, **526**, 225 - 230 (2020): "8-Nitro-cGMP Modulates Exocytosis in Adrenal Chromaffin Cells"

Kishimoto, Y.; Kunieda, K.; Kitamura, A.; Kakihana, Y.; Akaike, T.; Ihara, H., *ACS Chem. Neurosci.*, **9**, 217 - 223 (2018): "8-Nitro-cGMP Attenuates the Interaction between SNARE Complex and Complexin through S-Guanylation of SNAP-25"

Arimoto, H.; Takahashi, D., *Handb. Exp. Pharmacol.*, **238**, 253 - 268 (2017): "8-Nitro-cGMP: A Novel Protein-Reactive cNMP and Its Emerging Roles in Autophagy"

Yoshitake, J.; Soeda, Y.; Ida, T.; Sumioka, A.; Yoshikawa, M.; Matsushita, K.; Akaike, T.; Takashima, A., *J. Biol. Chem.*, **291**, 22714 - 22720 (2016): "Modification of Tau by 8-Nitroguanosine 3',5'-Cyclic Monophosphate (8-Nitro-cGMP): EFFECTS OF NITRIC OXIDE-LINKED CHEMICAL MODIFICATION ON TAU AGGREGATION"

Ihara H.; Ahtesham A. K.; Ida T.; Kasamatsu S.; Kunieda K.; Okamoto T.; Sawa T.; Akaike T., *Nitric Oxide*, **25**, 169 - 175 (2011): "Methodological Proof of Immunochemistry for Specific Identification of 8-nitroguanosine 3',5'-cyclic Monophosphate Formed on Glia Cells"

Ahtesham A. K.; Sawa T.; Akaike T., *Amino Acids*, **41**, 123 - 130 (2011): "Protein Cysteine S-guanylation and Electrophilic Signal Transduction by Endogenous Nitro-nucleotides"

Tokutomi Y.; Kataoka K.; Yamamoto E.; Nakamura T.; Fukuda M.; Nako H.; Toyama K.; Dong Y.-F.; Ahmed K. A.; Sawa T.; Akaike T.; Kim-Mitsuyama S., *Brit. J. Pharmacol.*, **162**, 1884 - 1893 (2011): "Vascular Responses to 8-nitro-cyclic GMP in Non-diabetic and Diabetic Mice"

Kurauchi Y.; Hisatsune A.; Isohama Y.; Sawa T.; Akaike T.; Shudo K.; Katsuki H., *J. Neurochem.*, **116**, 323 - 333 (2011): "Midbrain Dopaminergic Neurons Utilize Nitric Oxide/cyclic GMP Signaling to Recruit ERK that Links Retinoic Acid Receptor Stimulation to Up-regulation of BDNF"

Okamoto T.; Khan S.; Oyama K.; Fujii S.; Sawa T.; Akaike T., *J. Clin. Biochem. Nutr.*, **46**, 14 - 19 (2010): "A New Paradigm for Antimicrobial Host Defense Mediated by a Nitrated Cyclic Nucleotide"

Fujii S.; Sawa T.; Ihara H.; Tong K. I.; Ida T.; Okamoto T.; Ahtesham A. K.; Ishima Y.; Motohashi H.; Yamamoto M.; Akaike T., *J. Biol. Chem.*, **285**, 23970 - 23984 (2010): "The Critical Role of Nitric Oxide Signaling, via Protein S-Guanylation and Nitrated cyclic GMP, in the Antioxidant Adaptive Response"

Akaike T.; Fujii S.; Sawa T.; Ihara H., *Nitric Oxide*, **23**, 166 - 174 (2010): "Cell Signaling Mediated by Nitrated Cyclic Guanine Nucleotide"

Zaki M. H.; Fujii S.; Okamoto T.; Islam S.; Khan S.; Ahmed K. A.; Sawa T.; Akaike T., *J. Immunol.*, **182**, 3746 - 3756 (2009): "Cytoprotective Function of Heme Oxygenase 1 Induced by a Nitrated Cyclic Nucleotide Formed During murine Salmonellosis"

Sawa T.; Zaki M. H.; Okamoto T.; Akuta T.; Tokutomi Y.; Kim-Mitsuyama S.; Ihara H.; Kobayashi A.; Yamamoto M.; Fujii S.; Arimoto H.; Akaike T., *Nature Chem. Biol.*, **3**, 727 - 735 (2007): "Protein S-guanylation by the Biological Signal 8-nitroguanosine 3',5'-cyclic Monophosphate"